

Invariant Stein domains in Stein symmetric spaces and a non-linear complex convexity theorem

Simon Gindikin^{*} and Bernhard Krötz[†]

Abstract

We prove a complex version of Kostant's non-linear convexity theorem. Applications to the construction of G -invariant Grauert tubes of Riemannian symmetric G/K spaces are given.

Introduction

Let $X = G/K$ be a semisimple non-compact Riemannian symmetric space. We may assume that G is semisimple with finite center and we write $G = NAK$ for an Iwasawa decomposition of G . By our assumption, G sits in its universal complexification $G_{\mathbb{C}}$ and so $X \subseteq X_{\mathbb{C}} := G_{\mathbb{C}}/K_{\mathbb{C}}$. Note that $X_{\mathbb{C}}$ is a Stein symmetric space. Observe that the group G does not act properly (i.e. with compact isotropy subgroups) on $G_{\mathbb{C}}/K_{\mathbb{C}}$ since $K_{\mathbb{C}}$ is not compact. Write U for the compact real form of $G_{\mathbb{C}}$. The maximal connected subdomain of $GUK_{\mathbb{C}}/K_{\mathbb{C}} \subseteq X_{\mathbb{C}}$ on which G acts properly and which contains X was introduced in [AkGi90]. It is given by

$$\Xi := G \exp(i\Omega) K_{\mathbb{C}} / K_{\mathbb{C}}$$

where Ω is a polyhedral convex domain in $\mathfrak{a} := \text{Lie}(A)$ defined by

$$\Omega := \{X \in \mathfrak{a} : (\forall \alpha \in \Sigma) \ |\alpha(X)| < \frac{\pi}{2}\}.$$

Here Σ denotes the restricted root system with respect to $\mathfrak{a}$.

One of our principal aims is to construct a broad class of G -invariant Stein subdomains in Ξ . Let us remind the reader that it is a notoriously difficult problem to verify that certain unions of orbits of non-compact groups are Stein. For example the problem posed in [AkGi90] whether Ξ is Stein was unsolved until the last year.

The Iwasawa decomposition on G cannot be holomorphically extended to the whole group $G_{\mathbb{C}}$; we have that $N_{\mathbb{C}}A_{\mathbb{C}}K_{\mathbb{C}} \subsetneq G_{\mathbb{C}}$ is an open Zariski dense subset. The *Iwasawa domain* Ξ_I

^{*} Supported in part by the NSF-grant DMS-0070816 and the MSRI

[†] Supported in part by the NSF-grant DMS-0097314 and the MSRI

is defined as the maximal connected G -invariant subdomain in $X_{\mathbb{C}}$ which contains X and is contained in $N_{\mathbb{C}}A_{\mathbb{C}}K_{\mathbb{C}}/K_{\mathbb{C}}$:

$$\Xi_I := \left(\bigcap_{g \in G} g(N_{\mathbb{C}}A_{\mathbb{C}}K_{\mathbb{C}}/K_{\mathbb{C}}) \right)_0,$$

where $(\cdot)_0$ refers to the connected component containing X . Since Ξ_I is the connected component of an open intersection of Stein domains, it is easy to see that Ξ_I is Stein (cf. proof of Theorem 3.4).

The domains Ξ and Ξ_I again became of recent interest. One has $\Xi = \Xi_I$. Here are the results in chronological order:

- $\Xi \subseteq \Xi_I$ for all classical groups G (cf. [KrSt01a]).
- $\Xi = \Xi_I$ for all classical groups G (cf. [GiMa01]).
- $\Xi_I \subseteq \Xi$ for all G (cf. [Ba01]).

In [KOS01] it will be shown that $\Xi \subseteq \Xi_I$ for the exceptional cases. Also it is announced in [Hu01] that $\Xi \subseteq \Xi_I$. This will then give us $\Xi = \Xi_I$ in the general case. In particular, Ξ is Stein. Also, in the preprint [BHH01] a complex-geometric proof of the Steinness of Ξ is given.

Geometrically, the domain Ξ is a rather complicated object. If G is a group of Hermitian type, then $\Xi = X \times \overline{X}$ (cf. [BHH01], [GiMa01] or [KrSt01b]). If $X = G/K$ is classical, then there exists a group of Hermitian type $S \supseteq G$ with maximal compact subgroup $U \supseteq K$ such that G/K is a real form of the Hermitian symmetric space S/U . In [BHH01] and [KrSt01b] a G -equivariant subdomain $\Xi_0 \subseteq \Xi$ was exhibited which is biholomorphic to S/U . Further one has $\Xi = \Xi_0 = S/U$ if and only if Σ is of type C_n or BC_n . In particular, if $\Sigma \neq C_n, BC_n$, then $S/U \subsetneq \Xi$ and the explicit geometric structure of Ξ is very intricate. For many exceptional spaces G/K the geometric structure is even more complicated.

The domain Ξ (in [Gi98] it is called *complex crown of X*) is universal in the sense that many analytical and geometrical constructions on the Riemannian symmetric space X extend to Ξ . In [KrSt01a] it was shown that the inclusion $\Xi \subseteq \Xi_I$ implies that all eigenfunctions on X for the algebra of G -invariant differential operators $\mathbb{D}(X)$ extend holomorphically to Ξ . In [GiMa01] the problem $\Xi \subseteq \Xi_I$ is included in a broad class of geometrical problems connected with Matsuki duality. These geometrical problems include in particular the problem of the parametrization of compact complex cycles in flag domains (cf. [Wo92]).

Let us now come to the contents of this paper. Our first main result is a complex version of Kostant's non-linear convexity theorem (cf. [Kos73]). Write $G \rightarrow A$, $g \mapsto a(g)$ for the middle projection in $G = NAK$.

Theorem. (Kostant) *Let $X \in \mathfrak{a}$. Then*

$$a(K \exp(X)) = \exp(\text{conv}(\mathcal{W}X)),$$

where $\mathcal{W} = N_K(\mathfrak{a})/Z_K(\mathfrak{a})$ denotes the Weyl group and $\text{conv}(\cdot)$ refers to the convex hull of $(\cdot)$.

One shows that the middle projection $a: G \rightarrow A$ holomorphically extends to

$$(1) \quad a: G \exp(i\Omega)K_{\mathbb{C}} \rightarrow A_{\mathbb{C}}.$$

Then our main result is:

Theorem A. (Complex Convexity Theorem) *For all $X \in \Omega$ we have that*

$$a(G \exp(iX)) \subseteq A \exp(i \text{conv}(\mathcal{W}X)).$$

Our convexity theorem features interesting applications to the geometry of the domain Ξ and its generic subdomains which are defined as follows: Let $\omega \subseteq \Omega$ be a non-empty convex $\mathcal{W}$ -invariant open subset. Then we can form the domains

$$\Xi(\omega) := G \exp(i\omega) K_{\mathbb{C}} / K_{\mathbb{C}}.$$

Note that $\Xi = \Xi(\Omega)$ and that the Iwasawa projection (1) naturally factors to a holomorphic mapping $a: \Xi \rightarrow A_{\mathbb{C}}$. Finally we define for every $g \in G$ the *horospherical tube* $T(g, \omega) \subseteq X_{\mathbb{C}}$ by

$$T(g, \omega) := g(N_{\mathbb{C}} A \exp(i\omega) K_{\mathbb{C}} / K_{\mathbb{C}}).$$

As an application of Theorem A we now obtain:

Theorem B. *Let ω be an open convex Weyl group invariant subset of Ω . Then the following assertions hold:*

- (i) $a(\Xi(\omega)) = A \exp(i\omega)$.
- (ii) $\Xi(\omega) = (\bigcap_{g \in G} T(g, \omega))_0$.
- (iii) *The domain $\Xi(\omega)$ is Stein.*

Let us emphasize that in the the case of $\Xi(\omega) = \Xi$, the inclusion $\Xi \subseteq \Xi_I$ means only the existence of an Iwasawa projection $a: \Xi \rightarrow A_{\mathbb{C}}$; (i) in Theorem B gives a more precise information on the image of this projection. Further (ii) in the above theorem is a much stronger statement than (iii); in particular, (ii) implies (iii) since all horospherical tubes $T(g, \omega)$ are Stein.

Theorem B can be considered as an analogue of Lassalle's results for compact symmetric spaces (cf. [La78]). We can interpret Theorem A and Theorem B as statements for G -orbits in $X_{\mathbb{C}}$ intersecting $A \exp(i\Omega) K_{\mathbb{C}} / K_{\mathbb{C}}$ which in the case of compact symmetric spaces G/K are true for arbitrary G -orbits (cf. [La78]).

It is our pleasure to thank the MSRI, Berkeley, for its hospitality during the *Integral geometry program* where this work was accomplished. We thank Dmitri Akhiezer for his careful screening of the manuscript and his worthy suggestions. Further we would like to thank Karl-Hermann Neeb for his kindness to proofread the paper and Laura Geatti for sharing her knowledge with us on the boundary of Ξ .

1. Notation

Let G be a connected semisimple Lie group sitting inside a complexification $G_{\mathbb{C}}$. We denote by $\mathfrak{g}$ and $\mathfrak{g}_{\mathbb{C}}$ the Lie algebras of G and $G_{\mathbb{C}}$, respectively. Let $K < G$ be a maximal compact subgroup and $\mathfrak{k}$ its Lie algebra. Denote by $\theta: G \rightarrow G$ a Cartan involution which has K as a fixed point set.

Let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ the Cartan decomposition attached to $\mathfrak{k}$. Take $\mathfrak{a} \subseteq \mathfrak{p}$ a maximal Abelian subspace and let $\Sigma = \Sigma(\mathfrak{g}, \mathfrak{a}) \subseteq \mathfrak{a}^*$ be the corresponding root system. Related to this root system is the root space decomposition according to the simultaneous eigenvalues of $\text{ad}(H)$, $H \in \mathfrak{a}$:

$$\mathfrak{g} = \mathfrak{a} \oplus \mathfrak{m} \oplus \bigoplus_{\alpha \in \Sigma} \mathfrak{g}^{\alpha},$$

here $\mathfrak{m} = \mathfrak{z}_{\mathfrak{k}}(\mathfrak{a})$ and $\mathfrak{g}^{\alpha} = \{X \in \mathfrak{g}: (\forall H \in \mathfrak{a}) [H, X] = \alpha(H)X\}$. For the choice of a positive system $\Sigma^+ \subseteq \Sigma$ one obtains the nilpotent Lie algebra $\mathfrak{n} = \bigoplus_{\alpha \in \Sigma^+} \mathfrak{g}^{\alpha}$. Then one has the Iwasawa decomposition on the Lie algebra level

$$\mathfrak{g} = \mathfrak{n} \oplus \mathfrak{a} \oplus \mathfrak{k}.$$

We write A , N for the analytic subgroups of G corresponding to $\mathfrak{a}$ and $\mathfrak{n}$. For these choices one has for G the Iwasawa decomposition, namely, the multiplication map

$$N \times A \times K \rightarrow G, \quad (n, a, k) \mapsto nak$$

In particular, every element $g \in G$ can be written uniquely as $g = n(g)a(g)\kappa(g)$ with each of the maps $\kappa(g) \in K$, $a(g) \in A$, $n(g) \in N$ depending analytically on $g \in G$. The last piece of structure theory we shall recall is the little Weyl group. We denote by $\mathcal{W} = N_K(\mathfrak{a})/Z_K(\mathfrak{a})$ the *Weyl group* of $\Sigma(\mathfrak{a}, \mathfrak{g})$.

Finally we define the domain

$$\Omega = \{X \in \mathfrak{a}: (\forall \alpha \in \Sigma) \ |\alpha(X)| < \frac{\pi}{2}\}.$$

Clearly Ω is convex and $\mathcal{W}$ -invariant.

2. The complex convexity theorem

Let us first give the relevant notation. Denote by $K_{\mathbb{C}}$, $A_{\mathbb{C}}$, $N_{\mathbb{C}}$ the complexifications of K , A and N . Then $N_{\mathbb{C}}A_{\mathbb{C}}K_{\mathbb{C}}$ is a proper Zariski-open, hence dense subset of $G_{\mathbb{C}}$. Throughout this paper we will assume that $\Xi \subseteq \Xi_I$ holds, i.e.,

$$G \exp(i\Omega) \subseteq N_{\mathbb{C}}A_{\mathbb{C}}K_{\mathbb{C}}$$

(see the introduction for proofs of this statement). We now set $T_{\Omega} := A \exp(i\Omega) \subseteq A_{\mathbb{C}}$. Then $GT_{\Omega} \subseteq N_{\mathbb{C}}A_{\mathbb{C}}K_{\mathbb{C}}$.

One can show that one has a well defined holomorphic middle projection $N_{\mathbb{C}}A_{\mathbb{C}}K_{\mathbb{C}} \rightarrow A_{\mathbb{C}}/(A_{\mathbb{C}} \cap K_{\mathbb{C}})$. Due to the simple connectedness of T_{Ω} , this projection restricted to GT_{Ω} lifts to $A_{\mathbb{C}}$ (cf. [KrSt01a, proof of Th. 1.8(iii)]) and we obtain an analytic mapping

$$G \times T_{\Omega} \rightarrow A_{\mathbb{C}}, \quad (g, a) \mapsto a(ga)$$

holomorphic in the second variable such that

$$ga \in N_{\mathbb{C}}a(ga)K_{\mathbb{C}}$$

holds.

If V is a vector space and $E \subseteq V$ is a subset, then we denote by $\text{conv}(E)$ the convex hull of E in V .

Theorem 2.1. (Complex Convexity Theorem) *Assume that G is classical. Then we have for all $X \in \Omega$ that*

$$a(G \exp(iX)) \subseteq A \exp(i \text{conv}(\mathcal{W}X)). \quad \blacksquare$$

The proof of Theorem 2.1 will be given in several steps.

Fix $a \in T_{\Omega}$ and consider the function

$$f_a: K \rightarrow \mathfrak{a}_{\mathbb{C}}, \quad k \mapsto \log a(ka).$$

Further we write $p_{\mathfrak{a}_{\mathbb{C}}}: \mathfrak{g}_{\mathbb{C}} \rightarrow \mathfrak{a}_{\mathbb{C}}$ for the projection along $\mathfrak{k}_{\mathbb{C}} + \mathfrak{n}_{\mathbb{C}}$. For $x \in GT_{\Omega}K_{\mathbb{C}} \subseteq N_{\mathbb{C}}A_{\mathbb{C}}K_{\mathbb{C}}$ we write $b(x) = n(x)a(x)$ for the *triangular part* of x .

Lemma 2.2. *For any $a \in T_\Omega$, $k \in K$ and $X \in \mathfrak{k}$ we have*

$$\left. \frac{d}{dt} \right|_{t=0} f_a(\exp(tX)k) = p_{\mathfrak{a}_\mathbb{C}}(\text{Ad}(b(ka))^{-1}X).$$

Proof. This result can be easily deduced from the known case for $a \in A$ by analytic continuation. However, for convenience for the reader, we briefly recall the proof. We have

$$\begin{aligned} \left. \frac{d}{dt} \right|_{t=0} f_a(\exp(tX)k) &= \left. \frac{d}{dt} \right|_{t=0} \log a(\exp(tX)ka) \\ &= \left. \frac{d}{dt} \right|_{t=0} \log a(\exp(tX)b(ka)) \\ &= \left. \frac{d}{dt} \right|_{t=0} [\log a(b(ka)^{-1} \exp(tX)b(ka)) + \log a(ka)] \\ &= \left. \frac{d}{dt} \right|_{t=0} \log a(\exp(t \text{Ad}(b(ka))^{-1}X)) = p_{\mathfrak{a}_\mathbb{C}}(\text{Ad}(b(ka))^{-1}X), \end{aligned}$$

proving the lemma. ■

Write $\mathfrak{g}_\mathbb{C}^\mathbb{R}$ for $\mathfrak{g}_\mathbb{C}$ considered as a real Lie algebra and $\kappa_\mathbb{R}$ for the Cartan-Killing form on $\mathfrak{g}_\mathbb{C}^\mathbb{R}$. Let κ be the Cartan-Killing form on $\mathfrak{g}$ and recall the following relation between κ and $\kappa_\mathbb{R}$:

$$(\forall X, X', Y, Y' \in \mathfrak{g}) \quad \kappa_\mathbb{R}(X + iY, X' + iY') = 2(\kappa(X, X') - \kappa(Y, Y')).$$

For every $\lambda \in (\mathfrak{g}_\mathbb{C}^\mathbb{R})^*$ we define $H_\lambda \in \mathfrak{g}_\mathbb{C}^\mathbb{R}$ by $\lambda(X) = \kappa_\mathbb{R}(X, H_\lambda)$ for all $X \in \mathfrak{g}_\mathbb{C}^\mathbb{R}$.

For every $\lambda \in (\mathfrak{a}_\mathbb{C}^\mathbb{R})^*$ and $a \in T_\Omega$ we now define the function

$$f_{a,\lambda}: K \rightarrow \mathbb{R}, \quad k \mapsto \lambda(f_a(k)).$$

Lemma 2.3. *Let $a \in T_\Omega$ and $k \in K$. Then the following assertions hold:*

- (i) *For all $X \in \mathfrak{k}$ one has $\left. \frac{d}{dt} \right|_{t=0} f_{a,\lambda}(\exp(tX)k) = \kappa_\mathbb{R}(X, \text{Ad}(n(ka))H_\lambda)$.*
- (ii) *We have $df_{a,\lambda}(k) = 0$ if and only if*

$$\mathfrak{k} \perp_{\kappa_\mathbb{R}} \text{Ad}(n(ka))H_\lambda.$$

Proof. (i) is immediate from Lemma 2.2 and the notations introduced from above. (ii) follows from (i). ■

Write $i\mathfrak{a}_\mathbb{R}^*$ for the subspace of $(\mathfrak{a}_\mathbb{C}^\mathbb{R})^*$ which vanishes on $\mathfrak{a}$. Our next goal is to determine the critical set of $f_{a,\lambda}$ for $\lambda \in i\mathfrak{a}_\mathbb{R}^*$.

We also denote by θ the holomorphic extension of the Cartan involution to $G_\mathbb{C}$. Further we write $G_\mathbb{C} \rightarrow G_\mathbb{C}$, $x \mapsto \bar{x}$ for the complex conjugation with respect to the real form G .

Lemma 2.4. *Suppose that $X \in \Omega$ is regular. Set $a = \exp(iX)$ and let $k \in K$. Then the following implication holds*

$$ka \in NA_\mathbb{C}K_\mathbb{C} \Rightarrow k \in N_K(\mathfrak{a}).$$

Proof. We write $ka = nbk'$ for $n \in N$, $b \in A_\mathbb{C}$ and $k' \in K_\mathbb{C}$. Then

$$ka^2k^{-1} = ka(\theta(ka)^{-1}) = nbk'(\theta(nbk'))^{-1} = nb^2\theta(n)^{-1}.$$

Set $x := ka^2k^{-1}$. Then

$$\bar{x} = ka^{-2}k^{-1} = x^{-1}.$$

Thus we obtain that

$$\bar{x} = n\bar{b}^2\theta(n)^{-1} = \theta(n)b^{-2}n^{-1}.$$

Therefore $x = \theta(n)\bar{b}^{-2}n^{-1}$ and so

$$x^2 = \theta(n)\bar{b}^{-2}n^{-1}nb^2\theta(n)^{-1} \in A_{\mathbb{C}}\theta(N_{\mathbb{C}}).$$

Now let $0 < t \leq 1$ and write $a_t := \exp(itX)$ and accordingly we define x_t . Then we obtain that

$$\log x_t^2 = \log(k \exp(i4tX)k^{-1}) = \text{Ad}(k)(i4tX) \in \mathfrak{a}_{\mathbb{C}} + \theta(\mathfrak{n}_{\mathbb{C}})$$

for small t and hence for all t by analyticity. But $\text{Ad}(k)(i4tX) \in i\mathfrak{p}$. Hence by the regularity of X we obtain that $k \in N_K(\mathfrak{a})$, as was to be shown. $\blacksquare$

Proposition 2.5. *Suppose that $X \in \Omega$ and set $a = \exp(iX)$. Let $\lambda \in i\mathfrak{a}_{\mathbb{R}}^*$ be such that $H_{\lambda} \in i\mathfrak{a}$ is regular. Then we have*

$$df_{a,\lambda}(k) = 0 \iff k \in N_K(\mathfrak{a}).$$

Proof. The implication “ $\Leftarrow$ ” follows immediately from Lemma 2.3 (ii).

“ $\Rightarrow$ ”: Suppose that $df_{a,\lambda}(k) = 0$. Then Lemma 2.3 (ii) implies that $\mathfrak{k} \perp_{\kappa_{\mathbb{R}}} \text{Ad}(n(ka))H_{\lambda}$, or equivalently $\text{Ad}(n(ka))H_{\lambda} \in i\mathfrak{g} + \mathfrak{p}$. Assume that $k \notin N_K(\mathfrak{a})$. By Lemma 2.4 we have $n(ka) = \exp(Y)$ for some $Y \in \mathfrak{n}_{\mathbb{C}} \setminus \mathfrak{n}$. Hence

$$\text{Ad}(n(ka))H_{\lambda} = H_{\lambda} + \underbrace{[Y, H_{\lambda}]}_{\in \mathfrak{n}_{\mathbb{C}}} + \frac{1}{2} \underbrace{[Y, [Y, H_{\lambda}]]}_{\in [\mathfrak{n}_{\mathbb{C}}, \mathfrak{n}_{\mathbb{C}}]} + \dots$$

Thus the fact that $H_{\lambda} \in i\mathfrak{a}$ is regular implies that $\text{Ad}(n(ka))H_{\lambda} \in i\mathfrak{a} + (\mathfrak{n}_{\mathbb{C}} \setminus i\mathfrak{n})$. But this contradicts $\text{Ad}(n(ka))H_{\lambda} \in i\mathfrak{g} + \mathfrak{p}$, proving our proposition. $\blacksquare$

Proof of Theorem 2.1. First we observe that it is sufficient to prove

$$a(K \exp(iX)) \subseteq A \exp(i \text{conv}(\mathcal{W}X))$$

for all $X \in \Omega$. By a simple density/continuity argument we may further assume that X is regular. Suppose that there exists a $k \in K$ such that $a(k \exp(iX)) \notin A \exp(i \text{conv}(\mathcal{W}X))$, or equivalently

$$\text{Im} \log a(k \exp(iX)) \notin \text{conv}(\mathcal{W}X).$$

Then we find a regular element $\lambda \in i\mathfrak{a}_{\mathbb{R}}^*$ such that

$$f_{\exp(iX),\lambda}(k) > \max_{Y \in i \text{conv}(\mathcal{W}X)} \lambda(Y).$$

But Proposition 2.5 implies that $f_{\exp(iX),\lambda}$ takes its maximum at an element $k \in N_K(\mathfrak{a})$. Hence $f_{\exp(iX),\lambda}(k) = \lambda(\text{Ad}(k)iX)$; a contradiction to our inequality from above. $\blacksquare$

3. Applications

Let now $\omega \subseteq \Omega$ be an open Weyl group invariant convex set. Then we define the domain

$$\Xi(\omega) := G \exp(i\omega) K_{\mathbb{C}} / K_{\mathbb{C}}.$$

If $\omega = \Omega$, then we set $\Xi := \Xi(\Omega)$. Write $\partial\Xi(\omega)$ for the topological boundary of $\Xi(\omega)$ in $G_{\mathbb{C}}/K_{\mathbb{C}}$. Note the following properties of $\Xi(\omega)$:

- $\Xi(\omega)$ is open in $G_{\mathbb{C}}/K_{\mathbb{C}}$ (cf. [AkGi90]).
- $\Xi(\omega)$ is connected and G -invariant.
- G acts properly on Ξ (cf. [AkGi90]).
- One has $G \exp(i\partial\omega) K_{\mathbb{C}} / K_{\mathbb{C}} \subseteq \partial\Xi(\omega)$. Moreover if $\overline{\omega} \subseteq \Omega$, then we have $\partial\Xi(\omega) = G \exp(i\partial\omega) K_{\mathbb{C}} / K_{\mathbb{C}}$ (cf. [KrSt01b, Prop. 4.1]).

From our discussions in the previous section it is clear that we have a holomorphic projection

$$a: \Xi_{\omega} \rightarrow A_{\mathbb{C}}, \quad x \mapsto a(x)$$

with $x \in N_{\mathbb{C}} a(x) K_{\mathbb{C}} / K_{\mathbb{C}}$ for all $x \in \Xi$. Finally we define the Abelian tube domain

$$T_{\omega} := A \exp(i\omega) \subseteq A_{\mathbb{C}}.$$

An immediate consequence of our complex convexity theorem then is:

Lemma 3.1. *We have*

$$a(\Xi(\omega)) = T_{\omega}. \quad \blacksquare$$

Remark 3.2. From Lemma 3.1 we obtain in particular that $a(\Xi) \subseteq T_{\Omega}$. It is interesting to observe that this inclusion for $G = \mathrm{Sp}(n, \mathbb{R})$ extends a result of Siegel.

Consider the vector space $V := \mathrm{Sym}(n, \mathbb{R})$ of real symmetric matrices with its subcone of positive definite matrices V_+ . Then we have the symmetric Siegel domain

$$S^+ := V + iV_+ \subseteq V_{\mathbb{C}}.$$

Recall that $S^+ \cong \mathrm{Sp}(n, \mathbb{R})/U(n)$.

If we write Δ_j for the j -th principal minor on $V_{\mathbb{C}}$, then Siegel's Lemma says

$$(3.1) \quad \Delta_j(z) \neq 0 \quad \text{for } z \in S^+$$

and all $1 \leq j \leq n$. Now consider the rational functions

$$\chi_j(z) = \frac{\Delta_j(z)}{\Delta_{j-1}(z)}$$

on $V_{\mathbb{C}}$. Then the inclusion $a(\Xi) \subseteq T_{\Omega}$ for $G = \mathrm{Sp}(n, \mathbb{R})$ implies

$$(3.2) \quad \mathrm{Im} \chi_j(z) > 0 \quad \text{for } z \in S^+$$

and for all j . Note that (3.2) implies (3.1). To see this one identifies S^+ with the symmetric space $\mathrm{Sp}(n, \mathbb{R})/U(n)$. Then with $S^- := \overline{S^+}$ one has a biholomorphism $\Xi \cong S^+ \times S^-$ (cf. [GiMa01] or [KrSt01b]). Realizing A as diagonal matrices in G , one then easily shows that $a(\Xi) \subseteq T_{\Omega}$ implies (3.2).

The example discussed above admits a natural generalization to all tube domains $V + iV_+$ associated to an Euclidean Jordan algebra V and cone V_+ . ■

For every $g \in G$ we define the *horospherical tube* associated to ω by

$$T(g, \omega) := g(N_{\mathbb{C}}T_{\omega}K_{\mathbb{C}}/K_{\mathbb{C}}) \subseteq G_{\mathbb{C}}/K_{\mathbb{C}}.$$

Note that $T(\mathbf{1}, \omega)$ is biholomorphic to $N_{\mathbb{C}} \times T_{\omega}$. In particular, all horospherical tubes $T(g, \omega)$ are Stein.

Theorem 3.3. *For any non-empty open convex $\mathcal{W}$ -invariant set $\omega \subseteq \Omega$ the domain $\Xi(\omega)$ is the connected component of the intersection of horospherical tubes:*

$$\Xi(\omega) = \left(\bigcap_{g \in G} T(g, \omega) \right)_0.$$

Proof. From Lemma 3.1 we obtain that

$$\Xi(\omega) \subseteq N_{\mathbb{C}}T_{\omega}K_{\mathbb{C}}/K_{\mathbb{C}} = a^{-1}(T(\mathbf{1}, \omega)).$$

Thus the fact that $\Xi(\omega)$ is G -invariant and connected implies that $\Xi(\omega) \subseteq \left(\bigcap_{g \in G} T(g, \omega) \right)_0$. On the other hand we obtain from $\Xi = \Xi_I$ (cf. the discussion in the introduction) and Lemma 3.1 that $\Xi = \left(\bigcap_{g \in G} T(g, \Omega) \right)_0$. From this and Lemma 3.1 we hence get

$$\left(\bigcap_{g \in G} T(g, \omega) \right)_0 \subseteq \Xi \cap T(\mathbf{1}, \omega) = \Xi(\omega),$$

completing the proof of the theorem. ■

An interesting application of Theorem 3.3 is the following:

Proposition 3.4. *Let $\omega \subseteq \Omega$ be an open convex $\mathcal{W}$ -invariant set. Then the intersection*

$$I(\omega) := \bigcap_{g \in G} T(g, \omega)$$

is open. In particular, $I(\omega)$ and every connected component of $I(\omega)$ is Stein. In particular, $\Xi(\omega)$ is Stein.

Proof. (following a suggestion of Dmitri Akhiezer) Since $G = KAN$ and since AN leaves $T(\mathbf{1}, \omega)$ invariant, we obtain

$$I(\omega) = \bigcap_{k \in K} T(k, \omega).$$

Hence $I(\omega)$ is an intersection of open sets over the “compact parameter space” K . In particular I is open. Since all horospherical tubes $T(k, \omega)$ are Stein, we obtain that I is Stein. With $I(\omega)$ all its connected components are Stein, concluding the proof of the theorem. ■

As a final application of the complex convexity theorem we prove a result on the characterization of the boundary of Ξ .

Proposition 3.5. *Let $\omega \subseteq \Omega$ be an open convex $\mathcal{W}$ -invariant set. Let $(z_n)_{n \in \mathbb{N}}$ be a sequence with $z_n \rightarrow z_0 \in \partial\Xi(\omega)$. Then*

$$(\operatorname{Im} \log a(z_n))_{n \in \mathbb{N}}$$

is a sequence in ω and every accumulation point of this sequence lies in $\partial\omega$.

Proof. Set $X_n := \operatorname{Im} \log a(z_n)$. By Lemma 3.1 we have $X_n \in \omega$. Since $\overline{\omega}$ is compact in $\mathfrak{a}$, we may assume that $X_n \rightarrow X_0$ with $X_0 \in \overline{\omega}$.

If $X_0 \notin \partial\omega$, then we find a convex Weyl group invariant open set ω_1 such that $\overline{\omega_1} \subseteq \omega$ and $X_n \in \omega_1$. Then Theorem 3.3 implies that $z_n \in \Xi(\omega_1)$ and so $z_0 \in \Xi(\omega_1)$. Now $\overline{\omega_1} \subseteq \Omega$ and so $\partial\Xi(\omega_1) = G \exp(i\partial\omega_1)K_{\mathbb{C}}/K_{\mathbb{C}}$. Thus $z_0 \in \Xi(\omega)$, contradicting the assumption $z_0 \in \partial\Xi(\omega)$. This concludes the proof of the proposition. ■

References

- [AkGi90] Akhiezer, D. N., and S. G. Gindikin, *On Stein extensions of real symmetric spaces*, Math. Ann. **286**, 1–12, 1990.
- [Ba01] Barchini, L., *Stein Extensions of Real Symmetric Spaces and the Geometry of the Flag Manifold*, preprint.
- [BHH01] Burns, D., S. Halverscheid, and R. Hind, *The Geometry of Grauert Tubes and Complexification of Symmetric Spaces*, preprint.
- [Gi98] Gindikin, S., *Tube domains in Stein symmetric spaces*, Positivity in Lie theory: open problems, 81–97, de Gruyter Exp. Math., **26**, de Gruyter, Berlin, 1998.
- [GiMa01] Gindikin, S., and T. Matsuki, *Stein Extensions of Riemann Symmetric Spaces and Dualities of Orbits on Flag Manifolds*, MSRI preprint 2001-028.
- [Hu01] Huckleberry, A., *On certain domains in cycle spaces of flag manifolds*, preprint.
- [Kos73] Kostant, B., *On Convexity, the Weyl Group and the Iwasawa Decomposition*, Ann. scient. Éc. Norm. Sup. 4^e série, t. **6**, 413–455, 1973.
- [KrSt01a] Krötz, B., and R. J. Stanton, *Holomorphic extensions of representations: (I) automorphic functions*, preprint.
- [KrSt01b] —, *Holomorphic extensions of representations: (II) geometry and harmonic analysis*, preprint.
- [KOS01] Krötz, B., M. Otto, and R. J. Stanton, *Complex crowns of Riemannian symmetric spaces – exercises for the exceptional cases*, in preparation.
- [La78] Lassalle, M., *Séries de Laurent des fonctions holomorphes dans la complexification d’un espace symétrique compact*, Ann. Sci. École Norm. Sup. (**4**) **11** (1978), no. **2**, 167–210.
- [Wo92] Wolf, J. A., *The Stein condition for cycle spaces of open orbits on complex flag manifolds*, Ann. of Math. (**2**) **136** (1992), no. **3**, 541–555.

Simon Gindikin
 Department of Mathematics
 Rutgers University
 New Brunswick, NJ 08903
 USA
 gindikin@math.rutgers.edu

Bernhard Krötz
 The Ohio State University
 Department of Mathematics
 231 West 18th Avenue
 Columbus, OH 43210-1174
 USA
 kroetz@math.ohio-state.edu